

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

CERTIFICATE OF TRANSLATION

Honourable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

I, JOHN CHARLES McGILLEY, B.A. M.I.T.I., Technical Translator, of c/o
Priory Translations Limited, 11, Magdalen Street, Colchester, Essex, England,
hereby state:

THAT I am well acquainted with the French and English languages.

THAT I translated the document identified as French Patent Application No.
02 10560 filed at the National Institute of Industrial Property on 26th August 2002,
from French into English;

THAT the attached English translation is a true and correct translation of
French Patent Application No. 02 10560.

to the best of my knowledge and belief; and

THAT all statements made of my own knowledge are true and that all
statements made on information and belief are believed to be true and further,
that these statements are made with the knowledge that wilful false statements
and the like are punishable by fine or imprisonment, or both, under Section 1001
of Title 18 of the United States Code





JOHN CHARLES MCGILLEY

FRENCH REPUBLIC

inpi

NATIONAL INSTITUTE
OF INDUSTRIAL PROPERTY

Patent of invention

Utility certificate

OFFICIAL COPY

The Director General of the National Institute of Industrial Property certifies that the document annexed hereto is the certified true copy of an application for title of industrial property registered at the Institute.

Drawn up in Paris, 19th June 2007

For the Director General
of the National Institute
of Industrial Property
The Head of the
Patent Department

Martine PLANCHE

INPI

National Institute
of Industrial Property

26 bis, rue de Saint Petersburg
75800 Paris Cedex 08
Tel: 01 53 04 53 04 Fax: 01 42 94 86 54

PATENT OF INVENTION

Utility certificate

Intellectual property code – Book VI

Request for grant 1/2

Important! The 2nd page must be completed.

This form must be completed clearly in black ink

For INPI use only

Delivery of documents Date 26th August 2002 Place 75 INPI Paris National Registration No. assigned by INPI 0210560 Date of filing assigned by INPI 26th August 2002		1. Name and address of applicant or representative to whom correspondence should be addressed Monsieur André BOURGOUIN BEAUFOR IPSEN - S.C.R.A.S. Direction de la Propriété Industrielle 24 rue Erlanger 75781 PARIS CEDEX 16	
Your references for this file (optional) RS Cas 329 - ER/MM			
Confirmation of a filing by fax <input type="checkbox"/> no. assigned by INPI to the fax			
2. NATURE OF APPLICATION		Tick one of the following 4 boxes	
Patent application		<input checked="" type="checkbox"/>	
Application for utility certificate		<input type="checkbox"/>	
Divisional application <i>Initial patent application or initial application for utility certificate</i>		<input type="checkbox"/> No. Date / / No. Date / /	
Conversion of a European patent application <i>Initial patent application</i>		<input type="checkbox"/> No. Date / /	
3. TITLE OF INVENTION (200 characters or spaces maximum) Heterocarpine, a protein of vegetable origin with anticancer properties			
4. DECLARATION OF PRIORITY OR REQUEST FOR BENEFIT FROM THE FILING DATE OF A PREVIOUS FRENCH APPLICATION		Country or organisation Date: / / No. Country or organisation Date: / / No. Country or organisation Date: / / No. <input type="checkbox"/> If there are other priorities, tick the box and use the printed form "Continuation"	
5. APPLICANT		<input type="checkbox"/> If there are other applicants, tick the box and use the printed form "Continuation"	
Name or title of company		SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATIONS SCIENTIFIQUES (S.C.R.A.S.)	
First names			
Legal form		Simplified joint stock company	
SIREN no.		308197185	
APE-NAF code		741J	
Address	Street	42 rue du Docteur Blanche	
	Post code and town	75016 PARIS	
Country		FRANCE	
Nationality		French	
Telephone no. (optional)		(33) 01 44 30 43 43	
Fax no. (optional)		(33) 01 44 30 43 21	
Email address (optional)			

INPI

National Institute
of Industrial Property

For INPI use only

PATENT OF INVENTION

Utility certificate
Request for grant 2/2

Presentation of documents Date 26th August 2002 Place 75 INPI Paris National Registration no. assigned by INPI 0210560	
Your references for this file: (optional)	RS Cas 329 - ER/MM
6. REPRESENTATIVE	
Name	BOURGOUIN
First name	André
Office or company	BEAUFOR IPSEN - S.C.R.A.S. Direction de la Propriété Industrielle
No. of permanent power of attorney and/or contractual relationship	PG 8225
Address	Street 24 rue Erlanger
	Post code and town 75781 PARIS CEDEX 16
Telephone no. (optional)	(33) 01 44 96 10 10
Fax no. (optional)	(33) 01 44 96 13 42
Email address (optional)	andre.bourgouin@beaufour-ipsen.com
7. INVENTOR(S)	
The inventors are the applicants	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No In this case, supply a separate designation of inventor(s)
8. SEARCH REPORT	
Immediate establishment or deferred establishment	<input checked="" type="checkbox"/> <input type="checkbox"/>
Payment of fee in instalments	Payment in two instalments, only for natural persons <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No
9. REDUCTION IN LEVEL OF FEES	
	Only for natural persons <input type="checkbox"/> Requested for the first time for this invention (attach a notice of no charge) <input type="checkbox"/> Requested prior to this filing (attach a copy of the decision of acceptance for this invention or indicate its reference no.):
If you have used the printed form "Continuation", indicate the number of pages attached	
10. SIGNATURE OF APPLICANT OR REPRESENTATIVE: (name and capacity of signatory) André BOURGOIUN, Representative	Stamp of prefecture or INPI C. TRAN

The law no.78-17 of 6th January 1978 relating to data processing, files and freedoms applies to the replies given on this form. It guarantees the right of access and rectification for the data held about you with the INPI.

Heterocarpine, a protein of vegetable origin with anticancer properties

The present invention relates to protein of vegetable origin with anticancer properties which binds human GHRH (human Growth Hormone releasing hormone) .

5 Growth hormone ("GH") is a protein with 191 amino acids which stimulates the production of a number of growth factors, such as *Insulin-Like Growth Factor I* (IGF-1) and triggers the growth of a large number of tissues (skeleton, connective tissues, muscles and viscera). GH also has physiological activities, increasing the synthesis of the nucleic acids, proteins and lipolysis whilst reducing urinary secretions (Frohmman L.A. & Kineman, R.D., *Handbook of Physiology*, Hormonal Control of Growth, published by Kostyo, J.L. & Goodman, H.M. (Oxford Univ. Press, New York, 1999), p.
10 189-221).

The synthesis of GH is regulated by factors with positive or negative action secreted by the hypothalamus. The main factor controlling the production of GH is the "Growth Hormone Releasing Hormone" (GHRH), a peptide with 44 amino acids in humans.

15 GH and GHRH are involved in a number of diseases. Of these, the following should be mentioned in particular: cancer (in particular prostate or lung cancer), acromegaly, diabetic retinopathies and nephropathies; for these pathologies, treatment with GHRH antagonists is indicated. Due to the number of diseases potentially concerned, the industry continues to research GHRH antagonists.

20 The Applicant has therefore just isolated a new protein of vegetable origin, which has the property of binding human GHRH.

A first subject of the invention is therefore an isolated protein which can be obtained by extraction from the plant *Pilocarpus heterophyllus*, which is characterized in that it has a molecular mass of approximately 90.9 kDa and comprises fragments of peptide sequences SEQ.ID.NO.1, SEQ.ID.NO.2 and SEQ.ID.NO.3, said protein being able to
25 be presented in a glycosylated or non-glycosylated form. In order to simplify the disclosure which follows, this protein is hereafter called "heterocarpine".

Said SEQ.ID.NO.1, SEQ.ID.NO.2 and SEQ.ID.NO.3 sequences are as follows:

SEQ.ID.NO.1: KLGARYFDK

SEQ.ID.NO.2: YGEDIIVGVIDSGV

SEQ.ID.NO.3: PESESY

- 5 The nomenclature used above (as in the remainder of the present Application) in order to define the peptides is that specified by the "IUPAC-IUB Commission on Biochemical Nomenclature" in which, in accordance with the standard representation, the N-terminal amino acid (amino group) appears on the left and the C-terminal amino acid (carboxyl group) appears on the right. The term "natural amino acid" indicates one of the natural
- 10 L-amino acids found in the natural proteins: Gly, Ala, Val, Leu, Ile, Ser, Thr, Lys, Arg, Asp, Asn, Glu, Gln, Cys, Met, Phe, Tyr, Pro, Trp and His.

A protein is called "isolated" if it is taken out of its original environment. In particular, a natural protein is isolated if it is separated from the biological material with which it coexists in the natural system.

- 15 The invention preferably relates to heterocarpine in its non-glycosylated form.

According to a preferred variant of the invention, the heterocarpine is obtained from an extract of cells of the plant *Pilocarpus Heterophyllus* cultured *in vitro*.

Moreover, a subject of the invention is also a monoclonal antibody, or an antigen binding fragment of the latter, which specifically binds heterocarpine.

- 20 Heterocarpine has the property of binding human GHRH. *In vitro*, heterocarpine binds human GHRH and thus inhibits the synthesis of cyclic AMP induced during the binding of human GHRH on its receptor. *In vivo*, in rats, the heterocarpine/human GHRH complex is formed in the blood compartment and, in dose-dependent manner inhibits the GH synthesis induced by 10 µg of human GHRH in a mole to mole ratio. The
- 25 heterocarpine has the property of binding the human GHRH.

- These properties render the compounds of the invention suitable for a pharmaceutical use. Therefore, a subject of the invention is also, as a medicament, heterocarpine in glycosylated or non-glycosylated form. It also relates to pharmaceutical compositions containing, as active ingredient, heterocarpine in a glycosylated or non-glycosylated
- 30 form, said composition also comprising one or more pharmaceutically acceptable excipients. A further subject is the use of heterocarpine in a glycosylated or non-glycosylated form for preparing medicaments intended to antagonize the effects of

GHRH, to treat proliferative diseases (and in particular cancer), to treat acromegaly or to treat diabetic retinopathies and nephropathies. With regard to cancer, heterocarpine is particularly suitable for preparing a medicament intended to treat carcinoid and pancreatic tumors, hypothalamo-hypophyseal gangliocytomas, bronchial, intestinal and hepatic carcinomas, sympathoadrenergic tumors, pheochromocytomas, hypophyseal adenomas and thyroid carcinomas. Heterocarpine is particularly suitable for preparing a medicament intended to treat cancers the growth of which is dependent on the growth factor GHRH, and in particular for preparing a medicament intended to treat a cancer chosen from small-cell lung cancer and breast cancer (and quite particularly small-cell lung cancer).

A subject of the invention is also, as a medicament, a monoclonal antibody, or an antigen binding fragment of the latter, which specifically binds heterocarpine. It also relates to a pharmaceutical composition comprising, as active ingredient, a monoclonal antibody, or an antigen binding fragment of the latter, which specifically binds heterocarpine, said composition also comprising one or more pharmaceutically acceptable excipients. It relates moreover to the use of a monoclonal antibody, or of an antigen binding fragment of the latter, which specifically binds heterocarpine, for preparing medicaments intended to antagonize the effects of GHRH, to treat proliferative diseases (and in particular cancer), to treat acromegaly or to treat diabetic retinopathies and nephropathies. As regards cancer, said monoclonal antibody or said antigen binding fragment of the latter is particularly suitable for preparing a medicament intended to treat carcinoid and pancreatic tumors, hypothalamo-hypophyseal gangliocytomas, bronchial, intestinal and hepatic carcinomas, sympathoadrenergic tumors, pheochromocytomas, hypophyseal adenomas and thyroid carcinomas.

The invention also relates to the use of heterocarpine as an excipient in a pharmaceutical composition intended for the controlled release of GHRH. It also relates to a pharmaceutical composition comprising GHRH, heterocarpine and one or more pharmaceutically acceptable excipients.

Finally, other subjects of the invention are the processes making it possible to extract and isolate heterocarpine from cells of the plant *Pilocarpus Heterophyllus*, said cells preferably originating from *in vitro* cultures. These processes essentially include a stage of extraction of the cells from the plant *Pilocarpus Heterophyllus* with water at a temperature of 0 to 50°C, and preferably 4 to 25°C, said extraction stage being followed by a filtration stage to separate the heterocarpine-rich filtrate from the *Pilocarpus Heterophyllus* cells and by one or more stages of separation of the heterocarpine from the other components extracted from the plant *Pilocarpus Heterophyllus*.

According to a first variant, these extraction and isolation processes essentially include the following successive stages:

- 5 a) a stage of extraction of the cells from the plant *Pilocarpus Heterophyllus* with water at a temperature of 0 to 50°C, and preferably of 4 to 25°C, said extraction stage being followed by a filtration stage to separate the heterocarpine-rich filtrate from the *Pilocarpus Heterophyllus* cells;
- b) a stage of precipitation of the proteins extracted, for example by adding ammonium sulphate, followed by a stage of separation of the precipitate (by filtration or, preferably, by centrifugation);
- 10 c) solubilizing the precipitate recovered in Stage b) in water; and
- d) a stage of gel-filtration chromatography in order to separate the heterocarpine from the other components of the solution.

According to another variant, these extraction and isolation processes essentially include the following successive stages:

- 15 a) a stage of extraction of the cells from the plant *Pilocarpus Heterophyllus* with water at a temperature of 0 to 50°C, and preferably 4 to 25°C, said extraction stage being followed by a filtration stage to separate the heterocarpine-rich filtrate from the *Pilocarpus Heterophyllus* cells;
- 20 b) a stage of delipidation of the solution obtained in a), acidified by the addition of a non-oxidizing acid (for example hydrochloric acid, sulphuric acid or phosphoric acid) at a pH preferably comprised between 2 and 4, using liquid-liquid extraction (preferably by using an organic solvent such as dichloromethane, heptane, hexane or cyclohexane);
- 25 c) a stage of elimination of the tannins by bringing the delipidated solution obtained in c) into contact with polyvinylpyrrolidone (or also nylon 66) followed by filtration on large-pore resin (preferably a polystyrene-based resin such as the resin Diaion[®] HP-20);
- 30 d) adjusting the filtrate obtained after Stage c) to alkaline pH (preferably between pH 9 and 11) by the addition of a base such as ammonium hydroxide, sodium hydroxide or potassium hydroxide;
- 35 e) one or more stages of filtration on anion-exchange resin, the eluent for this filtration stage or these filtration stages preferably being a buffer solution having a pH between 9 and 11 and optionally containing concentration gradients of a salt (such as for example sodium chloride or ammonium sulphate), in order to separate the heterocarpine from the other components of the solution; and

f) a desalination stage consisting of passing the solution obtained in Stage e) over a resin separating the constituents of a mixture on the basis of their molecular mass (such as the resin Sephadex[®] G25 or Superdex[®] 200 HR) and the elution of this mixture over said resin with water.

5 The pharmaceutical compositions containing a compound of the invention can be in solid form such as, for example, powders, pills, granules, tablets, liposomes, gelatin capsules or suppositories. The pills, tablets or gelatin capsules can be coated with a substance capable of protecting the composition from the action of gastric acid or the enzymes in the subject's stomach for a sufficient period of time to allow this
10 composition to pass undigested into the latter's small intestine. The compound can also be administered locally, for example to the actual site of a tumor. The compound can also be administered according to a sustained-release process (for example by using a sustained-release composition or a perfusion pump). Appropriate solid supports can be, for example, calcium phosphate, magnesium stearate, magnesium carbonate, talc,
15 sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidine and wax.

The pharmaceutical compositions containing a compound of the invention can also be presented in liquid form such as, for example, solutions, emulsions, suspensions or a sustained-release formulation. Appropriate liquid supports can be, for example, water,
20 organic solvents such as glycerol or glycols such as polyethylene glycol, as well as their mixtures, in varying proportions, in water.

The administration of a medicament according to the invention can be carried out by topical, oral, parenteral route, by intramuscular injection etc.

25 The dose of a compound according to the present invention, to be provided for the treatment of the above-mentioned diseases or disorders, varies depending on the administration method, the age and body weight of the subject to be treated as well as the state of the latter, and will be finally decided by the attending doctor or vet. Such a quantity determined by the attending doctor or vet is here called "therapeutically effective quantity".

30 In accordance with the invention, the heterocarpine can be prepared by the process described hereafter.

Preparation of heterocarpine

According to a preferred variant of the invention, *in vitro* cultures of calluses or of cell suspensions originating from different organs of the plant have been carried out. These tissues cultured on semi-solid or liquid medium are able to bio-synthesize compounds having biological properties.

- 5 By "callus" is meant in the present Application a macroscopic cluster of undifferentiated cells of plants in culture on a semi-solid nutritive medium. The term "undifferentiated cells" in the present Application designates cells which have an aptitude under certain conditions to multiply in the form of a callus or of a cell suspension without any morphogenesis phenomenon. Finally, by "cell suspension", is
10 meant undifferentiated cells which can form microscopic clusters in culture in a liquid nutrition medium.

The choice of the nutritive medium, hormones, culture conditions forms an integral part of the invention as well as the extraction and analysis of the extracts from these *in vitro* cultures.

- 15 Cells from *Pilocarpus Heterophyllus* seeds can be cultured in suspension for example according to the procedure hereafter.

The organs are decontaminated according to the usual methods before being cultured. Plantule organs *in vitro* have also served as callogenesis starting material without requiring disinfection beforehand. The preferred basic nutritive medium is one of the
20 media commonly used for *in vitro* culture: this is Gamborg's medium (described in Gamborg et al., Nutrient requirements of suspension cultures of Soybean root cells, *Exp. Cell Res.* (1968), **50**(1), 151-158). The carbon source is saccharose but glucose can also be used at a concentration of 1 to 120 g/l, preferably approximately 30 g/l. The macro-elements content can also be reduced by a factor of 2. Auxin or an auxin and a
25 cytokinin are added to the medium, with a preference for a combination of both hormones, generally 2,4-dichlorophenoxyacetic acid and kinetin, but α -naphthaleneacetic acid (NAA), β -indoleacetic acid (IAA), β -indolbutanoic acid (IBA) or picloram can also be combined with the kinetin or benzylaminopurine (BAP). The concentration can vary from 0.1 to 10 mg/l for the auxin (for example 1 mg/l can be
30 chosen), and from 0.01 to 2 mg/l for the cytokinin (for example 0.06 mg/l can be chosen). The vitamins are those associated with the different basic media. The cultures are carried out in light or in darkness. The temperature can vary from 10°C to 33°C but is preferentially approximately 23°C. The pH of the medium is comprised between 4

and 6.5 and is preferentially adjusted to 5.8 before sterilization. Moreover, agar may or may not be added to the medium.

5 The primary calluses appear after a few days of culture and can be separated from the original implant, removed and subcultured after approximately 1 month then cultured on agar semi-solid medium (in tubes or Petri dishes), at intervals of 4 to 8 weeks, preferably 6 weeks, thus a callus can be kept for years by successive subcultures on new media. The callus can also be subcultured in a stirred liquid culture medium (Erlenmeyer flask or bioreactor) with subcultures at 2 to 6 weeks, preferably 3 weeks.

10 The strains obtained are distinguished by their genetic origin, culture conditions, appearance and absence of morphogenesis.

15 The lyophilized *Pilocarpus Heterophyllus* cells are extracted with water at a temperature of 0 to 50°C, and preferably 4 to 25°C. The extract thus obtained is lyophilized before being redissolved at a suitable concentration (for example approximately 30 % of dried matter). The proteins precipitated by the addition of a concentrated solution of ammonium sulphate (for example at a concentration representing 70 to 90 % of the saturation concentration) are dissolved in a minimum of water and the insoluble materials are recovered by centrifugation. The proteins are then separated by column chromatography (the eluent preferably being water) and the heterocarpine (identifiable by its molecular mass of approximately 90.9 kDa) can then
20 be recovered.

Preparation of antibodies specifically binding heterocarpine

25 The present invention provides binding agents, such as the antibodies which specifically bind heterocarpine. Such an agent is referred to as "specifically binding" a protein if it reacts at a detectable level (for example by an ELISA test) with said protein and does not detectably react with other proteins. "The binding" refers to a non-covalent association between 2 separate molecules such that a complex is formed. The binding ability can be evaluated, for example, by determination of the binding constant for the formation of the complex. The binding constant is the value obtained when the value of the complex concentration is divided by the product of the values of the non-complexed component concentration. 2 products are called "bound" when the binding constant
30 reaches 10³ l/mol. The binding constant can be determined using methods well known to a person skilled in the art.

Any agent which can satisfy the above criteria can be regarded as a binding agent.

In the present invention, a binding agent is preferably an antibody or a fragment of the latter. The antibody can be prepared by any technique available to a person skilled in the art (cf. Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988). In general, the antibodies can be produced by cell culture techniques including the generation of monoclonal antibodies or *via* transfections of antibody genes into host cells from bacteria or mammals in order to produce recombinant antibodies.

Among other techniques, use of those described hereafter is preferred. An immunogen containing heterocarpine is injected into a group of mammals (for example mice, rats, rabbits, sheep or goats). In this stage, the heterocarpine can serve as immunogen without modification. Alternatively, a superior immune response can be induced if the heterocarpine is combined with a transport protein such as bovine serum albumin or limpet haemocyanin. The immunogen is injected into the host animal, preferably according to a predetermined schedule, and the animals are bled periodically. Polyclonal antibodies specific to heterocarpine can thus be purified from such antiserum, for example, by affinity chromatography using heterocarpine coupled with an appropriate solid support.

Pharmaceutical compositions intended for the release of GHRH:

These compositions can in particular be prepared from heterocarpine and GHRH according to one of the methods described in the journal by De Wolf and Brett, *Pharmacological Reviews* (2000), **52**, 207-236 and the references cited therein.

Unless otherwise specified, all the technical and scientific terms used here have the same meaning as that usually understood by an ordinary specialist in the field to which this invention belongs. Similarly, all the publications, patent applications, all the patents and all other references mentioned here are incorporated by way of reference.

The following examples are presented in order to illustrate the above procedures and should in no event be considered as a limit to the scope of the invention.

OBTAINING THE HETEROCARPINE

Example 1:

Culture of cells in vitro:

A *Pilocarpus Heterophyllus* seed is germinated and the stem resulting from this germination is removed. Said stem is cultured in a Gamborg medium (Gamborg et al., Nutrient requirements of suspension cultures of Soybean root cells, *Exp. Cell Res.* (1968), 50(1), 151-158) to which 30 g/l of saccharose, 1 mg/l of 2,4-dichlorophenoxyacetic acid and 0.06 mg/l of kinetin have been added. The culture is carried out in tubes at a temperature of 23°C and in darkness. Subcultures are carried out every 6 weeks under usual conditions. The strains, which are granular in appearance, have a beige pigmentation.

A growth kinetic of the strains, based on the increase in mass of fresh and dried material from the biomass, was carried out over 8 weeks. The calluses from 2 tubes are combined and constitute a twice-weekly harvest, the first harvest taking place at time 0. Calluses and gelose are then harvested and lyophilized. It is observed that growth is exponential up to 6 weeks of culture before the appearance of a stationary growth phase.

15 *Extraction of the cell cultures:*

25 g of lyophilized *Pilocarpus Heterophyllus* cells are extracted twice by immersion in 375 ml of water at 4°C, and left overnight at 4°C, then in 250 ml of water at 4°C for 4 hours and finally washed with 125 ml of water at 4°C. Each aqueous solution thus obtained is filtered under vacuum through a glass filter surmounted with celite in order to separate the cell debris from the aqueous solution. The aqueous solutions thus combined are then lyophilized in order to obtain 9.4 g of dried material. The lyophilized dry extract is then dissolved in 31 ml of water at 20°C in order to obtain a solution containing 30 % of dry extract. 17.4 g of ammonium sulphate is added in small portions with constant magnetic stirring in order to precipitate the protein fraction. The protein precipitate is then separated from the ammonium sulphate solution by centrifugation at 3000 rpm for 20 minutes. The ammonium sulphate solution is decanted and the precipitated proteins are dissolved in 22 ml of water, re-centrifuged and filtered in order to eliminate the insoluble particles.

The filtrate obtained is then subjected to gel-filtration chromatography. It is injected into a column (Buchi N° 19678, L = 230 mm; internal diameter = 26 mm) filled with Superdex TM 200 (Amersham Pharmacia Biotech, reference no. 17-1043-01; particles with an average diameter of 13 µm) prepared according to the manufacturer's recommendations using ultra-pure water (Water's Milli-Q) as eluent at a flow rate of 5 ml per minute. 40 ml fractions are thus collected and the active protein is found in the third and fourth fraction. These fractions are lyophilized in order to obtain approximately 14.2 mg of active product.

The purity of the product obtained is demonstrated by the appearance of a single band on electrophoresis gel containing sodium dodecylsulphate (SDS PAGE). The product corresponding to this band is designated hereafter as heterocarpine.

Example 2:

- 5 The cells cultured *in vitro* according to the same procedure as that described in Example 1 above are extracted according to the method described hereafter.

100 g of lyophilized *Pilocarpus Heterophyllus* cells are extracted using 2 litres of demineralized water at 20°C, the mixture being maintained under stirring overnight. The cells and the extract are filtered by suction on frit (porosity 3, diameter 20 cm) covered by a celite bed (previously washed with acid; 1 to 2 cm thick). The cells recovered are washed with 400 ml of demineralized water before being eliminated. The aqueous filtrate is then acidified to pH 3.0 by the addition of approximately 10 ml of 18% hydrochloric acid. The acidified solution is then delipidated by liquid-liquid extraction using 400 ml of dichloromethane. The dichloromethane phase is decanted then eliminated. The delipidated solution is subjected to rotary evaporation in order to eliminate the residual dichloromethane. Approximately 30 g of polyvinylpyrrolidone is then added to the delipidated solution (pH approximately 3.0) and the mixture is stirred for approximately 30 minutes in order to eliminate the tannins. The mixture is filtered through a bed by suction on frit (porosity 3, diameter 10 cm) covered by a mixed bed comprising 25 g of celite (previously washed with acid) and 25 g of polyvinylpyrrolidone. The filtrate is then passed through a bed of 400 ml of Diaion[®] HP-20 (Mitsubishi Chemical Company) pre-activated according to the manufacturer's instructions. The resulting filtrate is then rendered alkaline (pH 10) by the addition of approximately 60 ml of a 20% ammonium hydroxide solution. A slight precipitation appears after 30 minutes of rest. 1 g of celite (previously washed with acid) is added to the alkaline solution which is then filtered by suction through a membrane filter (0.22 µm). Approximately 2 litres of filtrate are then passed through a HiPrep[®] Q XL 16/10 column, mounted on an Akta[®] purificator and pre-balanced at pH 10.2 with a piperazine/HCl 0.1M buffer, with a flow rate of 0.5 ml per minute (the HiPrep[®] column and the Akta[®] purificator are both products from the company Amersham Biosciences). The column is then washed successively with 6 column volumes of the starting buffer at pH 10.2, 5 column volumes of the same buffer containing a 0.2M concentration of NaCl; and 10 column volumes of the same buffer containing a 1M concentration of NaCl. The majority of the heterocarpine is recovered in the first 3 column volumes of buffer containing the 1M concentration of NaCl. The active fractions are desalinated by passing through a Sephadex[®] G25 column (volume of the bed: 260 ml) using demineralized water as eluent. The active fractions, found in the first column volume

corresponding to the hold-up volume, are then lyophilized in order to obtain 170 mg of heterocarpine. The heterocarpine thus obtained is practically single band on SDS PAGE gel.

CHARACTERIZATION OF THE HETEROCARPINE

Analysis and micro-sequencing:

- 5 The samples are loaded onto a 10 % polyacrylamide gel. After migration, the gels are fixed and stained with Coomassie blue.

10 The gel tracks represented in **Figure 3** corresponding to tracks 1, 2, 3, 4 and 5 are respectively the molecular weight marker (Amersham), 0.5, 1 and 2 µg of the content of the final heterocarpine fraction as obtained in Example 1 and the molecular mass marker (Amersham). Determination of the molecular mass by means of a standard molecular mass marker graph using standard computing tools well known to a person skilled in the art (for example, Viber Lourmat's Bio-Profil Bio1D software) makes it possible to show that heterocarpine has a molecular mass of 90.9 kiloDaltons (\pm 1.6 kiloDaltons).

- 15 For protein micro-sequencing analysis, the band of polyacrylamide containing the protein is cut out and digested in 300 µl of digestion buffer containing 50 mM Tris (pH 8.6), 0.03 % of sodium dodecylsulphate at 35°C for 18 hours in the presence of 0.4 µg of endolysine-C (Sigma). The peptides obtained are separated, by HPLC, on a DEAE-C18 in-line column 1 mm in diameter. The separation gradient is based on a mixture of acetonitrile (from 2 to 70 %) and 0.1 % trifluoroacetic acid (TFA). The sequencing is then carried out on a Procise sequencer (Applied Biosystem). In this way three peaks have been sequenced, making it possible to characterize heterocarpine in a unique manner. The corresponding sequences are identified in the present Application by SEQ.ID.NO.1, SEQ.ID.NO.2 and SEQ.ID.NO.3.

- 25 Analysis of the glycoproteins is carried out by the detection of sugared structures of the glycoproteins separated by SDS-PAGE gel. This detection system is a modification of the "Periodic Acid-Schiff" methods and leads to the appearance of magenta bands showing evidence of the glycoproteins (Sigma). For the heterocarpine as obtained in Example 1, the result reproduced in **Figure 4** is obtained.

30 **PHARMACOLOGICAL PROPERTIES OF HETEROCARPINE**

Stable transfections of the human GHRH receptor (hGHRH-R):

Human embryo kidney cells, HEK-293, (a cell line developed by Dr. Stuart Sealfon, Mount Sinai Medical School, New York, New York) expressing the human GHRH receptor in stable manner were obtained from Dr. Kelly Mayo (Northwestern
5 University, Chicago, IL).

Cell culture and membrane preparation:

The HEK-293 cells transfected in a stable manner with the human GHRH receptor described above are cultured in DMEM (Dulbecco's modified Eagle's medium, high glucose content; supplied by Life technologies) supplemented with 0.4 mg/ml of G418
10 (Life technologies) in the presence of 10 % of foetal calf serum and 4 mM of L-glutamine (Life technologies). The cells are homogenized in buffer A containing 50 mM HEPES (pH 7.4), 5 mM of magnesium chloride (MgCl₂), 2 mM of ethyleneglycol-bis(2-amino-ethyl)-N,N,N',N'-tetraacetic acid (EGTA) and 50 µg/ml of bacitracin then are subjected to sonication in the same buffer A. The cells thus homogenized are
15 centrifuged at 4°C at 39,000 g for 10 minutes, suspended in buffer A and re-centrifuged at 4°C at 40,000 g for 10 minutes. The total membrane proteins are quantified by Bradford's technique. The pelleted membranes are thus stored at -80°C for later use.

Competitive binding test on hGHRH-R:

The membranes of the HEK-293 cells transfected in a stable manner with the human
20 GHRH receptor are diluted to a concentration of 100 µg/ml in the reaction buffer containing 50 mM HEPES (pH 7.4), 5 mM of MgCl₂, 2 mM of EGTA, 50 µg/ml of bacitracin and 0.5 % of bovine serum albumin (BSA). The membranes are incubated with 0.05 nM of [¹²⁵I]GHRH(1-44 amide) (Amersham) in a final volume of 200 µl in the presence of increasing concentrations of heterocarpine for 2 hours at 23°C. The
25 reaction is stopped by rapid filtration on 96-well GF/C filters 0.1 % pre-loaded with polyethylenimine. The filters are then washed three times at 4°C with washing buffer containing 50 mM Tris (pH 7.4) using a Packard 96-well filtration station. The filters thus dried are submerged in 20 µl of scintillating cocktail (Microscint O, Packard) and are subjected to Topcount counting (Packard). The non-specific activity is determined
30 in the presence of 100 nM of hGHRH. A dose-response curve is generated for hGHRH (0.001 nM-100 nM) and the results obtained are included in **Figure 1**.

Competitive formation of cyclic AMP:

The HEK-293 cells transfected in a stable manner with the human GHRH receptor are distributed in 48-well culture plates and cultured for 3 days. The culture medium is then

removed and replaced by medium B containing 250 µl of DMEM (Dulbecco's modified Eagle's medium, high glucose content; supplied by Life technologies) in the presence of 0.5 % of BSA, 0.5 mM of 3-isobutyl-1-methylxanthine (IBMX) and pre-incubated for 5 minutes at 37°C. At the end of the pre-incubation period, the heterocarpine is tested for an additional 20 minutes. The concentrations observed are reported in **Figure 2**. The incubation is stopped by the addition of 100 µl of 0.1M HCl and the aliquots are analysed for their cyclic AMP content using the FlashPlate kit (New England Nuclear).

Assay of GH in rats:

Levels of GH in rats (Sprague Dawley males) are measured in blood samples by an enzymo-immunological test developed by Spi-Bio (Spi-Bio, France). The rats are treated by intravenous injection of heterocarpine at increasing doses (vehicle alone, 1, 3 and 10 nmol), then, 10 minutes after, by intravenous injection of 10 µg (3 nmol) of hGHRH. Ten minutes after injection of the hGHRH, the growth hormone levels are measured in the blood samples as described above. The results obtained are represented in **Figure 5**.

Measurement of anti-tumor activity :

Human tumor cells and in particular H-69 small-cell lung cancer cells are injected under the skin of athymic mice in order to produce a human tumor xenograft of approximately 80 mm³ some ten days after the first graft. The mice are treated every two days by intravenous injection of heterocarpine in increasing doses (vehicle alone, 2.5 mg/kg, 5 mg/kg and 10 mg/kg). The volume of the tumors is then measured every 4 days throughout the duration of the treatment.

Brief description of the figures:

Figure 1 is a graph representing the inhibition of binding of human GHRH on the human GHRH receptor as a function of increasing concentrations of heterocarpine.

Figure 2 is a graph representing the inhibition of the production of cyclic AMP in cells transfected in a stable manner with the human GHRH receptor in the presence of 10 nM of human GHRH as a function of increasing concentrations of heterocarpine.

Figure 3 is a reproduction of an SDS-PAGE protein gel plate showing the presence of heterocarpine having a molecular weight of 90.9 kDa.



Figure 4 is a reproduction of an SDS-PAGE protein gel plate showing that heterocarpine is a glycoprotein (Panel B).

Figure 5 is a representation in histogram form representing the inhibition of GH synthesis in rats in the presence of 10 μ g of human GHRH as a function of increasing concentrations of heterocarpine.

Claims

1. Use of an isolated protein which can be obtained by extraction from the plant *Pilocarpus heterophyllus*, said protein being characterized in that it has a molecular mass of approximately 90.9 kDa and comprises fragments of peptide sequences
5 SEQ.ID.NO.1, SEQ.ID.NO.2 and SEQ.ID.NO.3; said protein moreover being able to be presented in a glycosylated or non-glycosylated form, for preparing a medicament intended to treat cancers the growth of which is dependent on the growth factor GHRH.
2. Use according to claim 1, characterized in that the protein has been obtained from an extract from cells of the plant *Pilocarpus Heterophyllus* cultured *in vitro*.
- 10 3. Use according to claim 1 or 2, characterized in that the cancers the growth of which is dependent on the growth factor GHRH are chosen from small-cell lung cancer and breast cancer.
4. Use according to claim 3, characterized in that the cancer the growth of which is dependent on the growth factor GHRH is small-cell lung cancer.
- 15 5. Use according to claim 3, characterized in that the cancer the growth of which is dependent on the growth factor GHRH is breast cancer.

1/4

Figure 1Figure 2






Figure 3

**Figure 4**

**Figure 5**

SEQUENCE LISTING

<110> SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATION

<120> Heterocarpine, a protein of vegetable origin with anticancer properties

<130> RS 329 FR - sequence list

<140> FR 02/15560

<141> 2002-08-26

<160> 3

<170> PatentIn Ver. 2.1

<210> 1

<211> 10

<212> PRT

<213> Pilocarpus Heterophyllus

<400> 1
Lys Leu Ile Gly Ala Arg Tyr Phe Asp Lys
1 5 10

<210> 2

<211> 14

<212> PRT

<213> Pilocarpus Heterophyllus

<400> 2
Tyr Gly Glu Asp Ile Ile Val Gly Val Ile Asp Ser Gly Val
1 5 10

<210> 3

<211> 6

<212> PRT

<213> Pilocarpus Heterophyllus

<400> 3
Pro Glu Ser Glu Ser Tyr
1 5

INPI
National Institute of
Industrial Property

PATENT OF INVENTION
Utility certificate
Intellectual property code – Book VI

cerfa
No. 11235'02

Patent Department
26 bis, rue de Saint Petersburg
75800 Paris Cedex 08
Tel: 01 53 04 53 04 Fax: 01 42 93 59 30

Designation of inventor(s) Page no. 1/2
(If the applicant is not the inventor or
the sole inventor)

This form must be completed clearly in black ink

Your references for this file (optional)		RS 329 ER/MM	
National registration no.		02/10560	
Title of the invention (200 characters or spaces maximum)			
Heterocarpine, a protein of vegetable origin with anticancer properties			
Applicant(s): SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATIONS SCIENTIFIQUES (S.C.R.A.S.) 42 rue du Docteur Blanche 75016 PARIS FRANCE			
Designate(s) as inventor(s): (Indicate top right "Page no. 1/1". If there are more than three inventors, use an identical form and number each page, indicating the total number of pages).			
Name		FERRANDIS	
First names		Eric	
Address	Street	74 avenue Guy de Coubertin	
	Post code and town	78470 SAINT REMY LES CHEVREUSE	
Member company (optional)			
Name		TENG	
First names		Beng Poon	
Address	Street	6 chemin de la grange	
	Post code and town	91190 GIF-SUR-YVETTE	
Member company (optional)			
Name		SOHIER	
First names		Christine	
Address	Street	30 rue des Aubuis - Les Fourneaux	
	Post code and town	37390 SAINT ROCH	
Member company (optional)			
DATE AND SIGNATURE(S) OF APPLICANT(S) OR REPRESENTATIVE (Name and capacity of signatory) Paris, 12 September 2003			
André BOURGOUIN, Representative			

The law no. 78-17 of 6th January 1978 relating to data processing, files and freedoms applies to the replies given on this form. It guarantees the right of access and rectification for the data held about you with the INPI.

INPI
National Institute of
Industrial Property

PATENT OF INVENTION
Utility certificate
 Intellectual property code – Book VI

cerfa
 No. 11235'02

Patent Department
 26 bis, rue de Saint Petersburg
 7500 Paris Cedex 08
 Tel: 01 53 04 53 04 Fax: 01 42 93 59 30

Designation of inventor(s) Page no. 2/2
 (If the applicant is not the inventor or
 the sole inventor)

This form must be completed clearly in black ink

Your references for this file (optional)		RS 329 ER/MM
National registration no.		02/10560
Title of the invention (200 characters or spaces maximum)		
Heterocarpine, a protein of vegetable origin with anticancer properties		
Applicant(s): SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATIONS SCIENTIFIQUES (S.C.R.A.S.) 42 rue du Docteur Blanche 75016 PARIS FRANCE		
Designate(s) as inventor(s): (Indicate top right "Page no. 1/1". If there are more than three inventors, use an identical form and number each page, indicating the total number of pages).		
Name		THURIEAU
First names		Christophe
Address	Street	10 Bld Emile Augier
	Post code and town	75116 PARIS
Member company (optional)		
Name		
First names		
Address	Street	
	Post code and town	
Member company (optional)		
Name		
First names		
Address	Street	
	Post code and town	
Member company (optional)		
DATE AND SIGNATURE(S) OF APPLICANT(S) OR REPRESENTATIVE (Name and capacity of signatory) Paris, 12th September 2003		
André BOURGOUIN, Representative		

The law no. 78-17 of 6th January 1978 relating to data processing, files and freedoms applies to the replies given on this form. It guarantees the right of access and rectification for the data held about you with the INPI.